



Lay, A. C. (2021). Does FFAR4 agonism have therapeutic potential in cardiometabolic disease? *Endocrinology*, 162(10), [bqab145].  
<https://doi.org/10.1210/endocr/bqab145>

Peer reviewed version

License (if available):  
CC BY-NC-ND

Link to published version (if available):  
[10.1210/endocr/bqab145](https://doi.org/10.1210/endocr/bqab145)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Oxford University Press at <https://doi.org/10.1210/endocr/bqab145> . Please refer to any applicable terms of use of the publisher.

This journal has made an 'accepted manuscript' proof version openly available on the journal website.  
When the online version is updated from the AAM to the VOR, please download the VOR and add it to this record, then restrict the AAM

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

# Does FFAR4 agonism have therapeutic potential in cardiometabolic disease?

Abigail C Lay<sup>1</sup>

<sup>1</sup>Bristol Renal, Bristol Medical School, University of Bristol, UK, BS1 3NY

<https://orcid.org/0000-0001-5923-634X>

Corresponding author: Abigail Charlotte Lay, Email: mdacl@bristol.ac.uk

## KEY WORDS

FFAR4, GPR120, Diabetes, obesity, Metabolic syndrome, Cardiometabolic disease, vascular disease, atherosclerosis, CVD

## Commentary main text:

Metabolic conditions, such as diabetes and obesity, are major global health problems, with an increasing incidence and a significant burden on global healthcare. With diabetes and obesity comes a significantly higher risk of developing cardiovascular disease (CVD), which in turn is a major cause of morbidity and mortality among those individuals. As such, there is a pressing need to develop new treatment options which requires a better understanding of disease mechanisms and the identification of novel drug targets.

Fatty acids have a well-recognised role in metabolic diseases such as diabetes. As well as being a critical energy source, they act as ligands for Free Fatty acid receptors (FFARs); a group of G protein-coupled receptors, which mediate cellular signalling effects of fatty acids. One such receptor, FFAR4 (also known as GPR120), has been highlighted as essential in maintaining metabolic homeostasis; with insulin-sensitising and anti-inflammatory roles as well as controlling adipogenesis (1). As a receptor for long-chain fatty acids (LCFAs), FFAR4 has been shown to mediate some of the protective and beneficial cell signalling effects of omega-3 fatty acids (1,2).

In their article, Stuttgen and Sahoo (3) highlight the potential role of FFAR4 in the pathogenesis of cardiometabolic disease, noting that while there is extensive evidence of an involvement of FFAR4 in metabolic and inflammatory disease (including type 2 diabetes and obesity); the role of FFAR4 in the development of atherosclerosis and CVD is only recently emerging. Given the long-standing associations between FFAR4/GPR120 and obesity, diabetes and inflammation; and the considerable interest in developing therapeutic FFAR4 agonists (4), it is surprising that the role of FFAR4 in atherosclerosis and CVD (particularly in the context of metabolic disease) is not more well studied.

Their article (3) follows a recent publication which reports a cardioprotective role for FFAR4 in models of pressure overload and associations between the R270H polymorphism (affecting Gα/q signalling capacity) and eccentric remodelling in patient cohorts (5). Of note, this FFAR4 variant has previously been associated with an increased risk of obesity in European populations (6), though this association has not been consistently observed in different study populations (7) and the loss of FFAR4 expression or activity alone may not be sufficient to induce obesity. The important question also remains as to whether such FFAR4 variants increase the susceptibility to CVD in obese or diabetic individuals.

Another interesting consideration is whether FFAR4 activity could explain at least some of the cardioprotective effects of LCFAs (for example omega-3), similarly to the effects observed in diabetes, insulin resistance and inflammation (1,2), and whether variations in FFAR4 could account for some of the inconsistencies between association studies of LCFAs and CVD. Of course, LCFAs have multiple biological roles in addition to FFAR4 agonism, and further mechanistic studies are required to untangle the potential cardioprotective effects of LCFAs that are mediated by FFAR4.

Mechanistically, it seems likely that the involvement of FFAR4 in cardiometabolic disease is at least in part due to the role in modulating vascular inflammation. As detailed by Stuttgen and Sahoo (3) there are several lines of evidence supporting a role for FFAR4 in regulating macrophage activation; one of the beneficial effects of FFAR4 agonists reported in pre-clinical models is in improving chronic inflammation associated with insulin resistance, via regulating macrophage activity (4). This

potentially makes FFAR4 activation an attractive target for treating vascular damage, atherosclerosis and CVD, which are driven by chronic, low-grade inflammation. Indeed, there is also evidence that FFAR4 can limit monocyte-endothelial interactions (3). Though there is again some contradictory evidence for this role; in a study of FFAR4 in leukocytes, it was concluded that activation by polyunsaturated FAs had no effect in an animal model of atherosclerosis (8). In this instance it is important to note that the concentrations of FAs were comparatively much lower than those in other studies (1,2) and, crucially, the level of FFAR4 activation achieved by these FAs will not necessarily be comparable to the chronic and consistent FFAR4 activation possible with synthetic agonists. Further detailed studies using small-molecule FFAR4 agonists in pre-clinical models of atherosclerosis and CVD are therefore warranted.

Important sex-specific differences in the role of FFAR4 have also been noted (3), requiring further investigation and consideration in future studies. It will also be important to delineate the tissue- and cell-specific effects of FFAR4 activation, as the potential roles in chemoresistance and survival of certain cancers (3) suggests therapies based on activating FFAR4 may not always be beneficial.

As discussed by Stuttgen and Sahoo (3), there is a compelling case for a role of FFAR4 in cardiometabolic disease and there is clearly great potential for FFAR4 to be targeted therapeutically. Overall, pre-clinical studies investigating the effects of FFAR4 agonism in obesity, diabetes and inflammation are promising, but there is a distinct lack of research investigating the role of FFAR4 in atherosclerosis and CVD; leaving many important questions remaining. It seems variations in FFAR4 (including a loss of activity and expression) could play a role in susceptibility to cardiometabolic diseases and there is the potential that increasing FFAR4 activation could be protective; but further comprehensive studies, with a triangulation of findings from *in vitro* systems and pre-clinical models to larger population-based studies, are needed. Such future work will no doubt shed light on the mechanistic roles of FFAR4 in atherosclerosis and CVD and subsequently, whether FFAR4 agonism has broad therapeutic potential in cardiometabolic disease.

## DISCLOSURE STATEMENT

The author has nothing to disclose

## Funding details

ACL is currently a recipient of an Early Career Grant from The Society for Endocrinology, UK.

Accepted Manuscript

## REFERENCES

1. Hilgendorf KI, Johnson CT, Mezger A, Rice SL, Norris AM, Demeter J, Greenleaf WJ, Reiter JF, Kopinke D, Jackson PK. Omega-3 Fatty Acids Activate Ciliary FFAR4 to Control Adipogenesis. *Cell* 2019; 179:1289-1305.e1221
2. Hirasawa A, Tsumaya K, Awaji T, Katsuma S, Adachi T, Yamada M, Sugimoto Y, Miyazaki S, Tsujimoto G. Free fatty acids regulate gut incretin glucagon-like peptide-1 secretion through GPR120. *Nature medicine* 2005; 11:90-94
3. Stüttgen GM, Sahoo D. FFAR4: A New Player in Cardiometabolic Disease? *Endocrinology* 2021; 162
4. Oh DY, Walenta E, Akiyama TE, Lagakos WS, Lackey D, Pessentheiner AR, Sasik R, Hah N, Chi TJ, Cox JM, Powels MA, Di Salvo J, Sinz C, Watkins SM, Armando AM, Chung H, Evans RM, Quehenberger O, McNelis J, Bogner-Strauss JG, Olefsky JM. A Gpr120-selective agonist improves insulin resistance and chronic inflammation in obese mice. *Nature medicine* 2014; 20:942-947
5. Murphy KA, Harsch BA, Healy CL, Joshi SS, Huang S, Walker RE, Wagner BM, Ernste KM, Huang W, Block RC, Wright CD, Tintle N, Jensen BC, Wells QS, Shearer GC, O'Connell TD. Free fatty acid receptor 4 responds to endogenous fatty acids to protect the heart from pressure overload. *Cardiovascular research* 2021;
6. Ichimura A, Hirasawa A, Poulain-Godefroy O, Bonnefond A, Hara T, Yengo L, Kimura I, Leloire A, Liu N, Iida K, Choquet H, Besnard P, Lecoeur C, Vivequin S, Ayukawa K, Takeuchi M, Ozawa K, Tauber M, Maffei C, Morandi A, Buzzetti R, Elliott P, Pouta A, Jarvelin MR, Körner A, Kiess W, Pigeyre M, Caiazzo R, Van Hul W, Van Gaal L, Horber F, Balkau B, Lévy-Marchal C, Rouskas K, Kouvatsi A, Hebebrand J, Hinney A, Scherag A, Pattou F, Meyre D, Koshimizu TA, Wolowczuk I, Tsujimoto G, Froguel P. Dysfunction of lipid sensor GPR120 leads to obesity in both mouse and human. *Nature* 2012; 483:350-354
7. Vestmar MA, Andersson EA, Christensen CR, Hauge M, Glümer C, Linneberg A, Witte DR, Jørgensen ME, Christensen C, Brandslund I, Lauritzen T, Pedersen O, Holst B, Grarup N, Schwartz TW, Hansen T. Functional and genetic epidemiological characterisation of the FFAR4 (GPR120) p.R270H variant in the Danish population. *Journal of medical genetics* 2016; 53:616-623
8. Shewale SV, Brown AL, Bi X, Boudyguina E, Sawyer JK, Alexander-Miller MA, Parks JS. In vivo activation of leukocyte GPR120/FFAR4 by PUFAs has minimal impact on atherosclerosis in LDL receptor knockout mice. *Journal of lipid research* 2017; 58:236-246